

## PRODUCT INFORMATION LEAFLET

### 1. Product Name

**Brand Name:** Crocin 120

**Generic Name:** Paracetamol Paediatric Oral Suspension I.P.

### 2. Qualitative & Quantitative Composition

Each 5 ml contains:

Paracetamol I.P. 120 mg

In flavoured syrup base.

Colour: Carmoisine

### 3. Dosage Form

Oral Suspension

### 4. Clinical Particulars

#### 4.1. Indications/Uses

Analgesic and Antipyretic.

#### 4.2. Posology and method of administration

**Dosage:** 10-15 mg/kg per dose

Average Weight (Kg)	Dose* (ml)
10-12	6-7
12-14	7-8
14-18	8-11
18-20	11-12

\*Calculated based on 15 mg/kg body weight.

Maximum daily dosage is 60 mg/kg to be administered in divided doses of 10-15 mg/kg throughout the 24-hour period.

Do not take more frequently than every 4 hours. Always dose by body weight of child.

Do not take this medicine for more than 3 days without consulting doctor. Do not exceed the stated dose.

*For Children under 1 month:* Not Recommended.

*For children under 3 months:* If fever persists for more than 24 hours (4 doses) seek medical advice. This is to ensure that fever that may be due to a serious infection is quickly diagnosed.

*Use measuring cup available with the pack to measure the accurate dose.*

Use the smallest dose that you need to treat your symptoms and use the medicine for the shortest period of time necessary.

#### 4.3. Contra-indications

## CROCIN 120

Do not use Crocin 120 if child is allergic to paracetamol or any of the other ingredients in the product.

### 4.4. Warnings and Precautions

Crocin 120 contains Paracetamol. Do not take more than the recommended dose as it may cause serious harm to your liver. Do not use this medicine if you are taking any other prescription or non-prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu, or to aid sleep.

*Always read and follow the label*

In case of any of the below situations, you may need to avoid using this product altogether or limit the amount of paracetamol that you take.

#### **Check with your doctor before use if the child:**

- has liver or kidney problems.
- has depleted glutathione levels, such as those who have a severe infection, are severely malnourished, severely underweight as this may increase the risk of metabolic acidosis. Signs of metabolic acidosis include:
  - deep, rapid, difficult breathing,
  - feeling sick (nausea), being sick (vomiting),
  - loss of appetite).

Contact a doctor immediately if you get a combination of these symptoms. Please see your doctor if your symptoms do not improve.

Crocin 120 contains Methyl paraben and Propyl paraben. As parabens are derivative of benzoic acid, this may increase jaundice (yellowing of the skin and eyes) in babies.

### 4.5. Interaction with other medicaments and other forms of interaction

Before taking this medicine, make sure you consult your doctor if child is taking warfarin or similar medicines used to thin the blood.

### 4.6. Pregnancy and lactation

*Pregnancy:* As with the use of any medicine during pregnancy, pregnant women should seek medical advice before taking paracetamol. The lowest effective dose and shortest duration of treatment should be considered.

*Lactation:* Paracetamol is excreted in breast milk but not in a clinically significant amount at recommended dosages. Available published data do not contraindicate breastfeeding. Hence, can be taken during breastfeeding.

### 4.7. Effects on ability to drive and use machines, if contra-indicated

None

### 4.8. Undesirable effects/side effects

*Stop taking this medicine and tell your doctor immediately if:*

## CROCIN 120

- you experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- you experience a skin rash or peeling, or mouth ulcers.
- you have previously experienced breathing problems with aspirin or non-steroidal anti-inflammatory drugs, and experience a similar reaction with this product.
- you experience unexplained bruising or bleeding.

These reactions are rare.

### 4.9. Overdose

In case of overdosage, seek medical advice from a doctor immediately even if you do not have any symptoms because of the risk of liver failure.

In case of overdosage, you may also contact the **National Poisons Information Centre of India. Details of the same are as below:**

Department of Pharmacology  
All India Institute of Medical Sciences  
New Delhi-110029  
Toll Free No. - 1800 116 117  
Tel No.- 26589391, 26593677

## 5. Pharmacological Properties

### 5.1. Pharmacodynamic Properties & mechanism of action

**ATC code:** N02B E01

**Pharmacotherapeutic group:** Anilides

#### **Mechanism of Action**

Paracetamol is an analgesic and antipyretic. Its mechanism of action is believed to include inhibition of prostaglandin synthesis, primarily within the central nervous system.

#### **Pharmacodynamic Effects**

The lack of peripheral prostaglandin inhibition confers important pharmacological properties such as the maintenance of the protective prostaglandins within the gastrointestinal tract. Paracetamol is, therefore, particularly suitable for patients with a history of disease or patients taking concomitant medication in whom peripheral prostaglandin inhibition would be undesirable (such as, for example, those with a history of gastrointestinal bleeding or the elderly).

### 5.2. Pharmacokinetics

*Absorption:* Paracetamol is rapidly and almost completely absorbed from the gastrointestinal tract.

*Distribution:* Binding to the plasma proteins is minimal at therapeutic concentrations.

## **CROCIN 120**

*Metabolism:* Paracetamol is metabolised in the liver and excreted in the urine mainly as glucuronide and sulphate conjugates.

*Elimination:* Less than 5% is excreted as unmodified paracetamol. The mean plasma half life is about 2.3 hours.

### **6. Pharmaceutical Particulars**

#### **6.1. List of Excipients**

Sucrose  
Xanthan Gum  
Methyl Paraben  
Propyl Paraben  
Sodium Citrate  
Citric Acid Monohydrate  
Sorbitol Solution 70% NC  
Microcrystalline cellulose  
Sodium carboxy methyl cellulose  
Carmoisine  
Strawberry Flavour  
Purified Water

#### **6.2. Incompatibilities**

Not applicable

#### **6.3. Shelf life**

24 months

#### **6.4. Special storage conditions**

Keep out of sight and reach of children.

Store at ambient room temperature protected from light and moisture.

#### **6.5. Nature and specification of the container**

60 ml Amber colored PET bottles with ROPP cap along with a measuring cup.

#### **6.6. Instructions for Use and Handling**

Shake Well Before Use. Use measuring cup for correct dosing.

#### **6.7. Manufacturing License Holder**

Recipharm Pharmaservices Pvt. Ltd.

34<sup>th</sup> K.M., T-Begur, Tumkur Road, Nelamangala, Bangalore Rural- 562 123, India

#### **6.8. Marketed By**

GlaxoSmithKline Asia Private Limited,

Patiala Road, Nabha- 147201, Punjab, India

**\*Crocin is a trademark owned by or licensed to the GSK Group of companies**